

AD-A147 732

ELECTRONICALLY EXCITED ATOMIC AND MOLECULAR OXYGEN(U)
BRITISH COLUMBIA UNIV VANCOUVER DEPT OF CHEMISTRY
E A OGRYZLO 15 AUG 84 AFOSR-TR-84-0904 AFOSR-79-0088


1/1

UNCLASSIFIED

F/G 7/4

NL

END



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SEP 1984

(12)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 84-0904	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Electronically Excited Atomic and Molecular Oxygen		5. TYPE OF REPORT & PERIOD COVERED FINAL TECHNICAL REPORT - March 1, 1979 - June 30, 1984.
AUTHOR(s) ELMER A. OGRYZLO		6. PERFORMING ORG. REPORT NUMBER
PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry University of British Columbia Vancouver, B.C., CANADA, V6T 1Y6		8. CONTRACT OR GRANT NUMBER(s) AFOSR-79-0088
CONTROLLING OFFICE NAME AND ADDRESS AFOSR/NC Cdr: 410 Building AFB, DC. 20330-6448		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 61102F 2303/B1
MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 15, 1984
		13. NUMBER OF PAGES 15
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; Distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) same as 16.		
18. SUPPLEMENTARY NOTES B		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Electronic excitation, atmospheric airglow, atomic and molecular emission kinetics.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See reverse		

84 10 23 152

AD-A147 732

DTIC FILE COPY

DTIC
ELECTE
NOV 21 1984
S D B

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

↙
The rates and mechanisms through which the electronically excited states of oxygen are formed and destroyed in the upper atmosphere have been studied in a discharge flow system. Using a technique in which excited states are formed by surface catalyzed recombination, rate constants have been obtained for the quenching of the O₂(A), O₂(c) and O(singlet S) states by atmospheric species. The yields of the electronically excited states O₂(singlet delta), O₂(singlet sigma) and O(singlet S) in atom recombination have been determined directly. Evidence has been found for the existence of a precursor in each of these systems. The rate laws for the formation of O₂(A), O₂(c) and O(singlet S) have been established and rate constants for some of the elementary steps have been determined by combining these rate laws with the results of quenching experiments. Final report, AFOSR-79-0088, Electronically Excited Oxygen. ↘

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

AFOSR-TR. 84-0904

12

FINAL REPORT

ELECTRONICALLY EXCITED ATOMIC AND MOLECULAR OXYGEN

ELMER A. OGRYZLO

AFOSR-79-0088

UNIVERSITY OF BRITISH COLUMBIA

DEPARTMENT OF CHEMISTRY

2036 MAIN MALL

VANCOUVER, B.C. CANADA V6T-1Y6

1 March 1979 - 30 June 1984

84 10 152

Approved for public release;
distribution unlimited.

Final Report
Research Project 79-0088
Electronically Excited Atomic and Molecular Oxygen

Objectives:

The natural phenomenon whose understanding motivated this study is the night airglow. The objective of our study was the identification of the mechanism by which the observed emission from O_2 and O are excited at an altitude of about 100 km in the atmosphere of Earth and Venus.

On a more fundamental level our objective was to determine the rate and mechanism by which the recombination of oxygen atoms produces the electronically excited states of oxygen molecules and atoms.

Status of Research Effort:

The experiments undertaken in this project can be divided into two series. The first series of experiments attempted to produce a sudden high concentration of electronically excited oxygen by the surface catalyzed recombination of oxygen atoms on a nickel surface. These experiments were an attempt to determine the quenching and other relaxation processes of $O_2(A^3\Sigma^+)$, $O_2(A'^3\Delta)$, & $O_2(c^1\Sigma^-)$ which have not previously been determined. The second series attempted to determine the rate laws governing the formation of electronically excited states of O_2 in the recombination of oxygen atoms. The results of these experiments could then be combined with those from series 1 to calculate the absolute yield of each excited state in the recombination process.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFSC)
NOTICE OF TRANSMITTAL TO DTIC
This technical report has been reviewed and is
approved for public release IAW AFR 190-12.
Distribution is unlimited.
MATTHEW J. KERPER
Chief, Technical Information Division

Surface Catalyzed Recombination Studies

Attempts to produce an instantaneous concentrate of $O_2(A)$, $O_2(A')$, $O_2(c)$, and $O(^1S)$ met with success and rate constants for the quenching of these species by a number of gases were determined. These were reported in a series of papers (1,2,3,4,5) and are summarized in Table 1. Attempts to


TABLE 1: Rate constants for quenching of $O_2(A^3\Sigma_u^+)$, $O_2(A'^3\Delta_u)$ and $O_2(c^1\Sigma_u^-)$ by various gases

Quenching series	$O_2(A^3\Sigma_u^+)$ and $O_2(A'^3\Delta_u)$	$O_2(c^1\Sigma_u^-)$	$O(^1S)$
$O(^3P)$	$(1.29 \pm 0.1) \times 10^{-11} c, d$	$(5.9 \pm 0.6) \times 10^{-12}$	
$O_2(a^1\Delta_g)$	$(8.1 \pm 1.6) \times 10^{-11}$	$(6.0 \pm 2.4) \times 10^{-12}$	2×10^{-10}
O_2	1.3×10^{-13}	3×10^{-14}	
CO_2	7×10^{-13}	$< 6 \times 10^{-14}$	
SF_6	6×10^{-13}	2×10^{-13}	
N_2O	1.8×10^{-12}	4×10^{-13}	
He		6×10^{-15}	
Ar	7.2×10^{-16}	6×10^{-16}	

^c All units are $cm^3 \text{ molecule}^{-1} s^{-1}$

^d Indicated error limits are the 1 σ statistical uncertainties.

determine the temperature dependencies of these quenching processes were less successful. The only process for which we obtained data is the reaction:



Assuming that the temperature dependence can be represented by the Arrhenius equation we have plotted $\ln k_1$ versus $1/T$ in Figure 1.




Fig 1

The line drawn through these points yields:

$$k_1 = 2.9 \times 10^{-8} e^{- (1965/T)} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1} \quad (2)$$

Although k_1 was determined specifically for only the second vibrational level of the $\text{O}_2(\text{A})$ state we have reason to believe that it is not significantly different for any level which is formed on the nickel surface ($v' = 0$ to 5). Not only are the quenching rate constants for other levels insignificantly different but the distribution in these vibrational levels does not appear to change significantly with changing temperature.

The primary purpose of these experiments was to determine whether in the Earth's night airglow the quenching of the $\text{O}_2(\text{A})$ state by oxygen atoms is significant. Unfortunately the lowest temperature at which we could conduct our experiments was 232K. However, if we assume that the temperature dependence represented by equation 2 can be extended another 32°, then $k_1(200\text{K}) = 1.5 \times 10^{-12} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$. This is a factor of 20 smaller than the room temperature value. In the upper atmosphere where the peak oxygen atom concentration is $6.5 \times 10^{-11} \text{ atoms cm}^3$, the rate at which $\text{O}_2(\text{A})$ is quenched by O is about an order of magnitude smaller than the radiative decay rate (0.2 s^{-1}). We conclude, therefore, that oxygen atoms are not very important quenchers of the $\text{O}_2(\text{A})$ state in the upper atmosphere.

The magnitude of the preexponential in equation 2 calls for some comment. It is about two orders of magnitude greater than normal (i.e. that associated with gas kinetic collision frequency). Therefore it may not be

wise to assume that the temperature dependence of this rate constant is well represented by equation 2 much beyond the temperature range of these measurements, especially at high temperatures. This reservation does not affect our qualitative conclusion that oxygen atoms do not dominate the $O_2(A)$ removal rate in the upper atmosphere, since this would hold true even if the rate constant at 200K is no smaller than the value which we measured at 232K.

The measurement of rate constants for a series of vibrational levels of these excited states has not been possible except for the quenching of $O_2(A)$ by oxygen atoms where values were determined for the 5 lowest vibrational levels. These are summarized in Figure 2.




Fig. 2


Atom Recombination Studies

In these experiments the kinetic dependence of the emissions from the various excited states of oxygen that result from atom recombination were determined

1. Emission for $O(^1S)$ was found to obey the following rate law (6)

$$I = k[O]^2[M]/[O_2(a^1\Delta_g)]$$


where $k = 2.7 \times 10^{-27} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$. This rate law is consistent only with a "Barth Type" mechanism given by equations [1-5], where reaction [4] occurs at collision frequency $5 \times 10^{-10} \text{ cm}^3 \text{ molec}^{-1} \text{ s}^{-1}$ and k_1 is at least 25% of the total recombination rate constant ($4.75 \times 10^{-31} \text{ cm}^6 \text{ molec}^{-2} \text{ s}^{-1}$).



2. Emissions from $O_2(A^3\Sigma^+)$ and $O_2(A'^3\Sigma)$ were found to obey the rate law (7)

$$I = k[O]^2[M]/[O_2]$$

which is consistent with the simple mechanism given by reactions [6-8].






Combining these measurements with the quenching data provided by the surface catalyzed experiments yields a value for k_6 which is still almost an order of magnitude too small to explain the night airglow emission intensity from $O_2(A)$.

Emission from $O_2(c^1\Sigma^-)$ was found to obey the rate law (7)

$$I = [O][M]^2/[O_2]$$



Such a kinetic dependence suggests the presence of a precursor (O_2^*) involved in the process as described by reactions [9-13].



Quantum yields in these reactions have yet to be measured in the laboratory.

3. Emission from $O_2(b^1\Sigma^+)$ was found to have a complex kinetic dependence because of the unique involvement of ground state O_2 (8). Steady state and

time dependent measurements required the presence of a precursor (O_2^*) as indicated in the mechanism described by reactions [14-18]:



where O_2^* remains unidentified but $k_{15}/k_{17} = 2.6$ and $k_{14} > 1.6 \times 10^{34} \text{ cm}^6 \text{ molec}^{-2} \text{ s}^{-1}$.

4. Emission from $O_2(a^1\Delta_g)$ was found to have a complex kinetic dependence on molecular oxygen very similar to that observed for $O_2(b)$. The effect of adding molecular oxygen to a stream of oxygen atoms is shown in Figure 3.




FIGURE 3

The growth of $O_2(a)$ from atom recombination along a flow tube (time/cm = 6.4×10^{-3} s). The numbers next to each curve give the pressure of O_2 (in torr) added at the "15-cm position."

The rate constant in the absence of O_2 is less than 10% of the rate constant for total recombination. With O_2 present it rises by almost a factor of 10. The detailed kinetic analysis remains to be completed.

References

1. R.D. Kenner and E.A. Ogryzlo, *Int. J. Chem. Kinet.*, 7, 501 (1980).
2. R.D. Kenner and E.A. Ogryzlo, *J. Photochem.*, 18, 379 (1982).
3. R.D. Kenner and E.A. Ogryzlo, *Can. J. Chem.*, 61, 921 (1983).
4. R.D. Kenner and E.A. Ogryzlo, *Chem. Phys. Letts.*, 103, 209 (1984).
5. R.D. Kenner and E.A. Ogryzlo, *Can. J. Phys.*, December, 1984.
6. R.D. Kenner, E.A. Ogryzlo and P.T. Wassell, *Nature*, 291 (1981).
7. R.D. Kenner, E.A. Ogryzlo and S. Turley, *J. Photochem.* 10, 199, (1979).
8. E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell, *J. Photochem.*, 25, 389 (1984).

Publications Resulting from This Project

1. "On the Excitation of the Night Airglow on Earth, Venus and Mars", R.D. Kenner, E.A. Ogryzlo and S. Turley, J. Photochem., 10, 199 (1979).
2. "Deactivation of $O_2(A^3\Sigma^+)$ by O_2 , O and Ar", R.D. Kenner and E.A. Ogryzlo, Int. J. Chem. Kin., 7, 501 (1980).
3. "Excitation of the Green Line in the Night Airglow", R.D. Kenner, E.A. Ogryzlo and P.T. Wassell, Nature, 291, 398 (1981).
4. "A Direct Determination of the Rate Constant for the quenching of $O(^1S)$ by $O_2(a^1\Delta_g)$ ", R.D. Kenner and E.A. Ogryzlo, J. Photochem., 18, 379 (1982).
5. "Quenching of $O_2(c^1\Sigma^-)$ by $O(^3P)$, $O_2(a^1\Delta_g)$ and other Gases", R.D. Kenner and E.A. Ogryzlo, Can. J. Chem., 61, 921 (1983).
6. "Rate Constant for the Deactivation of $O_2(A^3\Sigma^+)$ by N_2 ", R.D. Kenner and E.A. Ogryzlo, Chem. Phys. Letts., 103, 209 (1983).
7. "Orange Chemiluminescence from Nitrogen Dioxide", R.D. Kenner and E.A. Ogryzlo, J. Chem. Phys., 80, 1 (1984).
8. "The Yield of $O_2(b^1\Sigma^+)$ in Oxygen Atom Recombination", E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell, J. Photochem., 25, 389 (1984).
9. "Quenching of the $O_2(A + X)$ Herzberg I Bands by $O_2(a^1\Delta_g)$ ", R.D. Kenner and E.A. Ogryzlo, Can. J. Phys., December (1984).

10. "Chemiluminescent Association Reactions in the Upper Atmosphere" by E.A. Ogryzlo in Chemiluminescence and Chemiionization, A. Fontijn Ed. Marcel Dekker, Amsterdam (1984).
11. "Chemiluminescence in Gas Phase Reactions" by R.D. Kenner and E.A. Ogryzlo in Chemiluminescence and Bioluminescence Today, J.G. Burr Ed. Marcel Dekker, Amsterdam (1985).

Professional Personnel Associated with Research Effort

Dr. R.D. Kenner (Post Doctoral Associate)

Dr. P.T. Wassell (Graduate student and Post Doctoral Fellow), Ph.D. 1981.

Interactions

Invited paper to be presented at the PAC conference, Honolulu, December 18, 1984. E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell, "The yield of $O_2(a^1\Delta_g)$ in Atom Recombination".

Invited Paper to be presented at I.G.U. Conference, San Francisco, December 2, 1984, E.A. Ogryzlo, "Laboratory Studies of the Excitation of O_2 in Atom Recombinations".

Invited Paper presented at the A.C.S. Conference, Philadelphia, August 24, 1984, E.A. Ogryzlo, "Chemiluminescence Association Reactions in the Upper Atmosphere".

Invited Paper presented at COSMO-84, Tampa, January 4, 1984, E.A. Ogryzlo, Y.Q. Shen and P.T. Wassell, "The Yield of $O_2(b^1\Sigma^+)$ in Oxygen Atom Recombination".

END

FILMED

12-84

DTIC